

MATHEMATICS

EXPONENTIAL SUMS DEVIATING LEAST FROM ZERO

BY

M. G. DE BRUIN AND H. VAN ROSSUM

(Communicated by Prof. N. G. DE BRUIN at the meeting of December 16, 1972)

Summary

In this paper exponential sums of the type $Y_1 e^{w_1 x} + Y_2 e^{w_2 x} + \dots + Y_n e^{w_n x}$ where w_1, w_2, \dots, w_n are pairwise distinct real numbers and Y_1, Y_2, \dots, Y_n are polynomials in a real variable x of degree $\mu_1, \mu_2, \dots, \mu_n$ (non-negative integers) respectively are considered. The problem posed is minimizing these sums in Tchebycheff norm on an interval $[-a, a]$ while satisfying the so called condition C (defined in section 1); the existence of a unique solution is then proved using the notion of a Tchebycheff-system (section 2).

At last convergence results for $\mu_1 + \mu_2 + \dots + \mu_n \rightarrow \infty$ are derived from results in the theory of the n -dimensional Padé-table for the functions $e^{w_1 x}, e^{w_2 x}, \dots, e^{w_n x}$ (section 3).

INTRODUCTION

In this paper exponential sums are considered as they arise in multi-dimensional Padé approximation. In a classical paper "Sur la généralisation des fractions continues algébriques" H. PADÉ [7] posed the following problem:

Determine polynomials X_1, X_2, \dots, X_n ($n \geq 2$) in the indeterminate x , not all identically zero, with complex coefficients and degrees at most $\mu_1, \mu_2, \dots, \mu_n$ (non-negative integers) respectively, such that

$$X_1 S_1 + X_2 S_2 + \dots + X_n S_n = 0(x^{\mu_1 + \mu_2 + \dots + \mu_n + n - 1})$$

where S_1, S_2, \dots, S_n are given power series in x all with non-zero constant term and where $0(x^{\mu_1 + \mu_2 + \dots + \mu_n + n - 1})$ denotes a power series in x in which the coefficients of $x^0, x^1, \dots, x^{\mu_1 + \mu_2 + \dots + \mu_n + n - 2}$ all vanish.

Under certain additional conditions the so-called Padé- n -tuple (X_1, X_2, \dots, X_n) is uniquely determined. It is placed in a cell with coordinates $(\mu_1, \mu_2, \dots, \mu_n)$. The configuration of all these cells forms the n -dimensional Padé-table for the power series (S_1, S_2, \dots, S_n) .

These n -dimensional Padé-tables did not receive much attention hitherto if $n > 2$ except for the treatment in the thesis of H. JAGER [5]. The case where $n = 2$ is equivalent with the well-known Padé-table of rational fractions approximating a given power series in the Padé sense. For a treatment of this case see O. PERRON [10], H. S. WALL [15]. In a few cases the n -dimensional Padé-table is known explicitly and we focus our attention on the case where $S_k = e^{w_k x}$ ($k = 1, 2, \dots, n$) with

$$w_1, w_2, \dots, w_n \in \mathbb{Q} \text{ and } w_k \neq w_l \text{ if } k \neq l \text{ (} k, l = 1, 2, \dots, n \text{)}.$$

In general an expression of the type $Y_1 e^{w_1 x} + Y_2 e^{w_2 x} + \dots + Y_n e^{w_n x}$ where the w_1, w_2, \dots, w_n are pairwise distinct and Y_1, Y_2, \dots, Y_n are polynomials in x of degree $\mu_1, \mu_2, \dots, \mu_n$ respectively is called an exponential sum of degree $\sum_{k=1}^n (\mu_k + 1) - 1$. These sums play an important rôle in non-linear approximation; here they appear in linear approximation problems.

We see that in n -dimensional Padé-approximation to the set of functions $\{e^{w_k x}\}_{k=1}^n$ special exponential sums arise deviating least from zero in the Padé sense, i.e. an approximation in the space of formal power series with "valuation metric". In section 1 we give a brief introduction to multi-dimensional Padé-approximation while in section 2 the notion of a multi-dimensional Walsh-array is introduced. This is a table analogous to the n -dimensional Padé-table but now the polynomials X_1, X_2, \dots, X_n are determined, satisfying condition C of the next page, in such a way that $X_1 S_1 + X_2 S_2 + \dots + X_n S_n$ is minimized in the Tchebycheff norm on a real interval; in our case the interval $[-a, a]$, $0 < a < \min \{\varrho_1, \varrho_2, \dots, \varrho_n\}$, where ϱ_i denotes the radius of convergence of S_i ($i = 1, 2, \dots, n$). In the case of multi-dimensional approximation to the set $\{e^{w_k x}\}_{k=1}^n$ we prove convergence properties of the n -dimensional Padé-table and the corresponding n -dimensional Walsh-array. This is done in section 3.

1. Let F denote the space of all formal power series in an indeterminate x (i.e. all series of the form $\sum_{n=m}^{\infty} a_n x^n$, m an integer and $a_m \neq 0$), with complex coefficients. Addition, scalar multiplication, multiplication and division are defined as usual.

First we define an absolute value $|\cdot|_F$ on F by:

$$|A|_F = e^{-m} \text{ when } A = \sum_{n=m}^{\infty} a_n x^n \text{ with } a_m \neq 0.$$

The aforementioned "valuation metric" ϱ is then defined as:

$$\varrho(A, B) = |A - B|_F, \quad A, B \in F.$$

Now let $S_k \in F$ with $|S_k|_F = 1$ ($k = 1, 2, \dots, n$) be n given elements of F . We ask for n polynomials (also elements of F) X_1, X_2, \dots, X_n of degree $\mu_1, \mu_2, \dots, \mu_n$ at most, respectively, such that

$$\left| \sum_{k=1}^n X_k S_k \right|_F \leq e^{-\mu_1 - \mu_2 - \dots - \mu_n - n + 1}.$$

From H. JÄGER [5], theorem 1.1.1, page 8, the existence of an n -tuple of polynomials like that follows immediately, while after specialisation to $S_k = e^{w_k x}$ ($k = 1, 2, \dots, n$; elements of F satisfying $|S_k|_F = 1$!) his theorems 1.2.1, 1.1.2 and 1.1.4 (page 13, 10 and 12 respectively) with $m = n$, $\varrho_k = \mu_k + 1$ ($k = 1, 2, \dots, n$) and $R = \mathbb{C}$ lead to:

Theorem 1.1. The problem of finding polynomials X_1, X_2, \dots, X_n

in x of degree $\mu_1, \mu_2, \dots, \mu_n$ at most respectively such that

$$\left| \sum_{k=1}^n X_k e^{w_k x} \right|_F \leq e^{-\mu_1 - \mu_2 - \dots - \mu_n - n + 1}$$

has a unique solution (apart from a multiplicative constant) which also satisfies:

- i) $\left| \sum_{k=1}^n X_k e^{w_k x} \right|_F = e^{-\mu_1 - \mu_2 - \dots - \mu_n - n + 1}$.
- ii) the degree of X_k is μ_k ($k = 1, 2, \dots, n$).

Remark: A set of power series admitting a unique solution (apart from a multiplicative constant) for the aforementioned problem satisfying i) and ii) is called *perfect*, a quite natural generalization of the notion normal in the case of the ordinary Padé-table.

Imposing one further condition on the coefficients of the polynomials appearing in the theorem we find a unique solution and place it in an n -dimensional table as described in the introduction.

The condition will be referred to as:

CONDITION C: the coefficient of x^{μ_1} in the polynomial X_1 is $1/\mu_1!$

Remark: In order to obtain a really unique solution it is sufficient either to fix one of the coefficients of the Padé-polynomials or to fix the first non-zero coefficient of $\sum_{k=1}^n X_k(\mu_1, \mu_2, \dots, \mu_n; x) e^{w_k x}$. Taking the last mentioned coefficient to be $\alpha/(\mu_1 + \mu_2 + \dots + \mu_n + n - 1)!$ ($\alpha \in \mathbb{C}$, "symmetric" condition), the coefficient of x^{μ_1} in X_1 will be $\alpha/\mu_1! \prod_{r=2}^n (w_1 - w_r)^{\mu_r + 1}$ and conversely.

Imposing a condition on the coefficient of the highest power of x appearing in X_1 can be seen as a sort of generalization of the condition in the well-known Tchebycheff problem: find the monic polynomial of degree n deviating least from zero on $[-1, 1]$ in Tchebycheff-norm.

The deeper reason for the chosen normalization lies in the possibility to derive a result on uniqueness and convergence in an n -dimensional Walsh-array from those in the corresponding n -dimensional Padé-table. Uniqueness is ascertained in the Walsh case by the requirement that we have a so called Tchebycheff-system and it is precisely the chosen normalization that leads to such a system in our case (cf. Lemma 3.2).

In the n -dimensional Padé-table the sequence of cells $(0, 0, \dots, 0)$, $(1, 1, \dots, 1)$, $(2, 2, \dots, 2)$, ..., $(\lambda, \lambda, \dots, \lambda)$, ... is called the *main diagonal* and the cells satisfying $\mu_1 + \mu_2 + \dots + \mu_n = k$ the *plane of approximation of order k*. Now a sequence of cells will be called *regular* if any two arbitrary consecutive cells are adjacent and if the second cell lies in a plane of approximation of order one higher than the order of the first one. The recurrence relations for Padé n -tuples in an n -dimensional Padé-table constitute a generalization of the continued fraction algorithms connected with regular sequences in the ordinary Padé-table. Research on these algorithms closely

related to the Jacobi-Perron algorithms (investigated by O. PERRON [8], [9] published thirteen years after H. PADÉ [7]) is carried on by the first mentioned author of this paper.

Now consider the two-dimensional case. Let $S_1, S_2 \in F$ with $|S_1|_F = |S_2|_F = 1$ and assume $(V_{\mu\nu}^{(1)}, V_{\mu\nu}^{(2)})$ to be the Padé-pair for the square (μ, ν) . Then $|S_1 V_{\mu\nu}^{(1)} + S_2 V_{\mu\nu}^{(2)}|_F = e^{-\mu-\nu-1}$ or equivalently $S_1 V_{\mu\nu}^{(1)} + S_2 V_{\mu\nu}^{(2)} = 0(x^{\mu+\nu+1})$. Dividing both members by $S_2(x)$ and putting $C(x) = S_1(x)/S_2(x)$ we obtain

$$C(x) V_{\mu\nu}^{(1)}(x) + V_{\mu\nu}^{(2)}(x) = 0(x^{\mu+\nu+1}).$$

So $P_{\mu\nu}(x) = -V_{\mu\nu}^{(2)}(x)/V_{\mu\nu}^{(1)}(x)$ is the Padé-approximant on the square (μ, ν) of the ordinary Padé-table for $C(x)$.

In the metric space F every sequence $\{P_{\mu\nu}(x)\}$ of Padé-approximants to $C(x)$ converges to $C(x)$ if $\mu + \nu \rightarrow \infty$. If x is an element of a certain subset of \mathbb{C} however the question of convergence (and uniform convergence) on this set is much more complicated.

For power series $C(x) = \sum_{n=0}^{\infty} c_n x^n$ where $\{c_n\}_{n=0}^{\infty}$ is a so-called *Stieltjes-sequence* fairly detailed results are available; see for instance G. A. BAKER Jr. etc. [1], H. S. WALL [15]. Also the work of A. F. BEARDON [2], [3] and J. S. R. CHISHOLM [4] should be mentioned.

Regarding the pointwise and uniform convergence for sequences $\{P_{\mu\nu}(x)\}$ in which μ and ν augment by unity alternately (so called *step-lines*, examples of regular sequences) the work of the first author of this paper (to appear) is of some interest. For all known functions with a normal table the Padé-approximants on the steplines appear to be approximants of continued fractions which converge to the function on the whole region of convergence of its Taylor series around the origin; this implies convergence of the sequence of Padé-approximants on the diagonals.

2. Consider expressions of the form $W_1 S_1 + W_2 S_2 + \dots + W_n S_n$ in which S_1, S_2, \dots, S_n are power series with real coefficients in the real variable x with radius of convergence $\varrho_i > 0$ ($i = 1, 2, \dots, n$) respectively and W_1, W_2, \dots, W_n polynomials with real coefficients of degree $\mu_1, \mu_2, \dots, \mu_n$ respectively; let $[-a, a]$ be a closed finite interval with $0 < a < \min\{\varrho_1, \varrho_2, \dots, \varrho_n\}$.

We now want to minimize the expression

$$\max_{|x| \leq a} |W_1 S_1 + W_2 S_2 + \dots + W_n S_n|$$

under condition C for the n -tuple W_1, W_2, \dots, W_n .

In order to be able to solve the problem and maybe even find a unique solution it must be reformulated and some conditions must be imposed upon S_1, S_2, \dots, S_n . These conditions are laid down in the concept of a *Tchebycheff-system* (see S. KARLIN, W. J. STUDDEN [6]), which can also be used in much more general "minimax" problems.

The problem can be reformulated as:

$$\text{minimize } \max_{|x| \leq a} \left| \frac{x^{\mu_1}}{\mu_1!} S_1 - (-W_1^* S_1 - W_2 S_2 - \dots - W_n S_n) \right|,$$

where degree $W_1^* = \mu_1 - 1$, degree $W_k = \mu_k$ ($k = 2, 3, \dots, n$).

(If $\mu_1 = 0$ we minimize $\max_{|x| \leq a} |S_1 - (-W_2 S_2 - \dots - W_n S_n)|$).

Definition 2.1. A system $\{u_0, u_1, \dots, u_m\}$ of continuous real-valued functions on a closed finite interval $[a, b]$ is called a *Tchebycheff-system* (*T-system*) when the determinants

$$\begin{vmatrix} u_0(t_0) & u_0(t_1) & \dots & u_0(t_m) \\ u_1(t_0) & u_1(t_1) & \dots & u_1(t_m) \\ \vdots & \vdots & & \vdots \\ u_m(t_0) & u_m(t_1) & \dots & u_m(t_m) \end{vmatrix}$$

are strictly positive whenever $a \leq t_0 < t_1 < \dots < t_m \leq b$.

Example: In the case $u_i(t) = t^i$ ($i = 0, 1, \dots, m$) the determinants reduce to familiar Vandermonde determinants from which the positiveness follows for an arbitrary finite interval.

It will be convenient for the sequel to use a slightly different way of ascertaining the *T-system* property:

Theorem 2.1. If a system $\{u_i\}_{i=0}^m$ of continuous real-valued functions on $[a, b]$ satisfies the condition that every non-trivial (i.e. $\neq 0$) function $\alpha_0 u_0 + \alpha_1 u_1 + \dots + \alpha_m u_m$ ($\alpha_i \in \mathbb{R}$; $i = 0, 1, \dots, m$) has at most m zeros on $[a, b]$ then $\{u_i\}_{i=0}^m$ is a *T-system* except possibly for the sign of one of the functions.

(For the proof see S. KARLIN, W. J. STUDDEN [6], theorem 4.1, page 22).

Theorem 2.2. The function $\alpha_0 u_0 + \alpha_1 u_1 + \dots + \alpha_m u_m$ minimizing

$$\max_{|x| \leq a} \left| f(x) - \sum_{i=0}^m \alpha_i u_i(x) \right|$$

where f is a continuous real-valued function on $[-a, a]$ is uniquely determined if and only if the functions $\{\varepsilon u_0, u_1, \dots, u_m\}$ constitute a *T-system* for appropriate $\varepsilon = +1$ or -1 .

(Proof: S. KARLIN, W. J. STUDDEN [6], theorem 1.1, page 280).

Bearing this in mind we now assume $\{S_1, xS_1, \dots, x^{\mu_1-1}S_1, S_2, \dots, x^{\mu_2}S_2, \dots, S_n, \dots, x^{\mu_n}S_n\}$ to be a *T-system* on $[-a, a]$. In this case the minimizing problem posed at the beginning of this section has a unique solution and placing this unique n -tuple in the cell $(\mu_1, \mu_2, \dots, \mu_n)$ of an n -dimensional array we arrive at what will be called an *n-dimensional Walsh-array*. For $n=2$, $S_2=1$, we arrive at a slightly different form of the ordinary Walsh-array ([17]). Convergence properties in the ordinary Walsh-array have been considered by E. B. SAFF [12], [13], [14] and J. L. WALSH [18].

3. Consider the functions $S_k = e^{w_k x}$ ($k=1, 2, \dots, n$); $w_1, w_2, \dots, w_n \in \mathbb{C}$ pairwise distinct and let G be a bounded subset of \mathbb{C} ; $\mu_1, \mu_2, \dots, \mu_n$ non-negative integers.

According to theorem 1.1 there exists a unique n -tuple of polynomials $P_k(\mu_1, \mu_2, \dots, \mu_n; x)$ ($k=1, 2, \dots, n$) satisfying condition C and:

- i) $|\sum_{k=1}^n P_k(\mu_1, \mu_2, \dots, \mu_n; x) e^{w_k x}|_F = e^{-\mu_1 - \mu_2 - \dots - \mu_n - n + 1}$
- ii) degree $P_k(\mu_1, \mu_2, \dots, \mu_n; x) = \mu_k$ ($k=1, 2, \dots, n$).

These polynomials and the sum in i) can be given explicitly using H. JÄGER [5], chapter IV, § 2 with $m=n$, $\rho_k = \mu_k + 1$ ($k=1, 2, \dots, n$). For the sequel we only need:

$$\begin{aligned} R(x) &= \sum_{k=1}^n P_k(\mu_1, \mu_2, \dots, \mu_n; x) e^{w_k x} = \\ &= \frac{\prod_{v=2}^n (w_1 - w_v)^{\mu_v + 1}}{2\pi i} \int_{|z|=R} \frac{e^{xz}}{\prod_{k=1}^n (z - w_k)^{\mu_k + 1}} \frac{dz}{z} \end{aligned}$$

where R is a fixed real number satisfying:

$$R > 2 \max [|w_1|, |w_2|, \dots, |w_n|, |w_1 - w_2|, |w_1 - w_3|, \dots, |w_1 - w_n|, 1].$$

We are now able to formulate the following theorem which will be proved by estimating $R(x)$:

Theorem 3.1.

$$\lim_{\mu_1 + \mu_2 + \dots + \mu_n \rightarrow \infty} \sum_{k=1}^n P_k(\mu_1, \mu_2, \dots, \mu_n; x) e^{w_k x} = 0$$

uniformly in x on G .

Proof: Put $\mu_1 + \mu_2 + \dots + \mu_n = \sigma$ and let M be a bound for G , i.e. for all $x \in G$: $|x| \leq M$.

Then:

$$\begin{aligned} |R(x)| &< \frac{(\max \{|w_1 - w_2|, |w_1 - w_3|, \dots, |w_1 - w_n|, 1\})^{\sigma + n - 1 - \mu_1}}{2\pi} \cdot \frac{2\pi R e^{MR}}{(\frac{1}{2}R)^{\sigma + n} \cdot R} < \\ &\leq d^{\sigma + n} \cdot e^{MR} \text{ where } d = \frac{\max \{|w_1 - w_2|, |w_1 - w_3|, \dots, |w_1 - w_n|, 1\}}{\frac{1}{2}R} \end{aligned}$$

satisfying $0 < d < 1$, which shows the uniform convergence for $\sigma \rightarrow \infty$.

Remark: This theorem still holds when we use the aforementioned condition on the first non-zero coefficient of $R(x)$ with $\alpha=1$ instead of condition C . The difference is that we have to omit the scalar $\prod_{v=2}^n (w_1 - w_v)^{\mu_v + 1}$ in the formula for $R(x)$ and we can omit $\max \{|w_1 - w_2|, |w_1 - w_3|, \dots, |w_1 - w_n|, 1\}$ in the lower bound for R and in the proof of the theorem.

We now want to formulate a minimax problem involving the functions $e^{w_k x}$ ($k=1, 2, \dots, n$) and therefore we restrict ourselves to the case $w_k \in \mathbb{R}$ ($k=1, 2, \dots, n$) and G a real interval $[-a, a]$, $a > 0$.

Lemma 3.2. The problem of minimizing

$$\max_{|x| \leq a} \left| 0 - \sum_{k=1}^n A_k(\mu_1, \mu_2, \dots, \mu_n; x) e^{w_k x} \right|$$

where the minimum is taken over all polynomial n -tuples with degree $A_k = \mu_k$ ($k=1, 2, \dots, n$) satisfying condition C has a unique solution; this solution will be referred to as $\{W_k(\mu_1, \mu_2, \dots, \mu_n; x)\}_{k=1}^n$.

Once this lemma is proven we arrive at the following important theorem:

Theorem 3.2.

$$\lim_{\mu_1 + \mu_2 + \dots + \mu_n \rightarrow \infty} \sum_{k=1}^n W_k(\mu_1, \mu_2, \dots, \mu_n; x) e^{w_k x} = 0$$

uniformly in x on $[-a, a]$.

Remark: If μ_1 is fixed the theorem still holds when $\sigma \rightarrow \infty$, showing the convergence to be a non-trivial matter.

Proof of theorem 3.2: Since the Padé polynomials belong to the set of n -tuples over which the minimum is taken, lemma 3.1 leads to:

$$\begin{aligned} \left| \sum_{k=1}^n W_k(\mu_1, \mu_2, \dots, \mu_n; x) e^{w_k x} \right| &\leq \max_{|x| \leq a} \left| \sum_{k=1}^n W_k(\mu_1, \mu_2, \dots, \mu_n; x) e^{w_k x} \right| < \\ &\leq \max_{|x| \leq a} \left| \sum_{k=1}^n P_k(\mu_1, \mu_2, \dots, \mu_n; x) e^{w_k x} \right| < d^{\sigma+n} \cdot e^{MR} \end{aligned}$$

($0 < d < 1$, M and R as in theorem 3.1); the last expression converging to zero for $\sigma \rightarrow \infty$.

Proof of lemma 3.2: Using the same method for reformulating the minimax problem as in section 2 we only have to prove that

$$\{e^{w_1 x}, x e^{w_1 x}, \dots, x^{\mu_1-1} e^{w_1 x}, e^{w_2 x}, \dots, x^{\mu_2} e^{w_2 x}, \dots, e^{w_n x}, \dots, x^{\mu_n} e^{w_n x}\}$$

is a T -system on $[-a, a]$ except possibly for the sign of one of the functions and then use theorem 2.2. (If $\mu_1 = 0$, consider the system $\{e^{w_2 x}, \dots, x^{\mu_2} e^{w_2 x}, \dots, e^{w_n x}, \dots, x^{\mu_n} e^{w_n x}\}$).

A linear combination with real coefficients of the $\sigma + n - 1$ functions of the system gives rise to an exponential sum of degree at most $\{(\mu_1 - 1) + 1 + \sum_{k=1}^{n-1} (\mu_k + 1)\} - 1 = \sigma + n - 2$. According to G. POLYA, G. SZEGO [11], page 48, problem 75 an exponential sum has at most as many real zeros as its order, so the linear combination of the $\sigma + n - 1$ functions has at most $\sigma + n - 2$ zeros on $[-a, a]$ and thus the functions form a T -system (except possibly for the sign of one of them) in view of theorem 2.1.

REFERENCES

1. BAKER JR, G. A., J. L. GAMMEL and J. G. WILLS, An investigation of the applicability of the Padé-approximant method. *Journ. Math. Anal. Appl.* 2, 405-418 (1961).
2. BEARDON, A. F., On the convergence of Padé-approximants. *Journ. Math. Anal. Appl.* 21, 344-346 (1968).
3. ———, On the location of the poles of Padé-approximants. *Journ. Math. Anal. Appl.* 21, 469-474 (1968).
4. CHISHOLM, J. S. R., Approximation by sequences of Padé-approximants in regions of meromorphy. *J. Math. Phys.* 7, 39-44 (1966).
5. JAGER, H., A multidimensional generalization of the Padé table. Thesis Amsterdam (1964).
6. KARLIN, S. and W. J. STUDDEN, *Tchebycheff systems with applications in analysis and statistics*. New York, London, Sydney (1966).
7. PADÉ, H., Sur la généralisation des fractions continues algebriques. *J. de Math. Pures et Appl.* X, 291-329 (1894).
8. PERRON, O., Grundlagen für eine Theorie des Jacobischen Kettenbruchalgorithmus. *Math. Ann.* 64, 1-76 (1907).
9. ———, Über die Konvergenz der Jacobi Kettenbruchalgorithmen mit Komplexen Elementen. *Sitzungsber. d. Bay. Ak. (math. Phys. Klasse)* 37, 401-482 (1907).
10. ———, *Die Lehre von den Kettenbrüchen II*. Stuttgart (1957).
11. POLYA, G. und G. SZEGO, *Aufgaben und Lehrsätze aus der Analysis II*. Springer, Berlin etc. (1964).
12. SAFF, E. B., On the row convergence of the Walsh-array for meromorphic functions. *Trans. Amer. Math. Soc.* 146, 241-257 (1969).
13. ———, The convergence of best approximation to the exponential function I. *Notices Amer. Math. Soc.* 71, 196 (1970).
14. ———, Idem II, *Notices Amer. Math. Soc.* 71, 1073 (1970).
15. WALL, H. S., *The analytic theory of continued fractions*. Toronto, New York, London (1948).
16. WALSH, J. L., Padé-approximants as limits of functions of best approximation. *J. Math. and Mech.* 13, 305-312 (1964).
17. ———, On approximation to an analytic function by rational functions of best approximation. *Math. Z.* 38, 163-176 (1934).
18. ———, The convergence of sequences of rational functions of best approximation with some free poles. *Approximation of functions* (Editor H. L. Garabedian), Elsevier Amsterdam (1965).